

Epidemiological characteristics of mortality from infectious intestinal diseases (diarrhea) Paraguay (2015-2019)

Características epidemiológicas de la mortalidad por enfermedades infecciosas intestinales (diarreas), Paraguay (2015 – 2019)

Eva Fabiana Mereles Aranda ^{1,2}, Juan Edgar Tullo Gómez ³, Derlis Duarte Zolán ^{1,3}, Elder Oliveira da Silva ⁴, Myrian Elizabeth Rivas Kiese ¹, Pasionaria Ramos ⁵.

¹Research Center in Health and Environmental Science (ScienHE), Asunción, Paraguay

²MICROBAC, Investigación y Microbiología, Ciudad del Este, Paraguay.

³Ministerio de Salud Pública y Bienestar Social, Asunción, Paraguay.

⁴Universidad de Salvador, Facultad de Medicina, Salvador, Brasil

⁵Universidad Nacional de Asunción, Facultad de Ciencias Médicas, Asunción, Paraguay



Received: 03/11/2023

Revised: 15/11/2023

Accepted: 24/12/2023

Corresponding author

Pasionaria Ramos. Universidad Nacional de Asunción, Paraguay
pasionariaramos@gmail.com

Responsible editor

Mg. Iván Barrios

Conflicts of interests

The authors declare that there is no conflict of interest.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

This article is published under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).



ABSTRACT

Introduction: Infectious intestinal diseases (diarrhea) are a major cause of morbidity and mortality worldwide. In 2015, it constituted the ninth leading cause of death for all ages. **Objective:** To describe the epidemiological characteristics of mortality due to infectious intestinal diseases (diarrhea) in Paraguay from 2015 to 2019. **Methods:** Descriptive, ecological, cross-sectional, retrospective, and non-probabilistic sampling of consecutive cases was performed. ICD-10 deaths included cholera (A01), typhoid and paratyphoid fever (A01), shigellosis (A03), other bacterial intestinal infections (A04), other bacterial food poisoning (A05), amebiasis (A06), other intestinal diseases due to protozoa (A07), intestinal infections due to viruses (A08), and diarrhea and gastroenteritis of presumed infectious origin (A09). Open access data were obtained from the WEB page of the General Directorate of Strategic Information in Health of the MSP and BS of all the Departments of the country. Crude and adjusted rates (per 100,000 inhabitants) were calculated. Excel and EPI INFO 7.0 were used. **Results:** 495 deaths were registered; the highest prevalence was in 2019 (adjusted rate of 1.83) and the lowest in 2017 (1.36). The months with the highest mortality rates were January and July (9.9%). 51.72% were women (256), 30.91% were over 80 years old and 28.28% (140) were under 5 years old, average age 50.8, single 56.77% (281) and without any type of education 41.82% (207). A total of 80.61% (399) of the respondents resided in urban areas. The highest rates were registered in Boquerón (33.3) and Amambay (15.2). Diarrhea and gastroenteritis of infectious origin accounted for 95.56% of the cases (473). **Discussion:** There was a high percentage of mortality from infectious diarrhea. Extremes of life were the most vulnerable populations.

Keywords: gastroenteritis; infections; diarrhea; Paraguay.

RESUMEN

Introducción: Las enfermedades infecciosas intestinales (diarrea) son una causa importante de morbilidad y mortalidad a nivel mundial. En 2015, constituyeron la novena causa principal de muerte para todas las edades. **Objetivo:** describir las características epidemiológicas de la mortalidad por enfermedades infecciosas intestinales (diarreas) en Paraguay durante los años 2015 al 2019. **Metodología:** Estudio descriptivo, ecológico, transversal, retrospectivo, muestreo no probabilístico de casos consecutivos. Se consideraron los óbitos del CIE – 10, que incluyen al cólera (A01), fiebres tifoidea y paratifoidea (A01), shigelosis (A03), otras infecciones intestinales bacterianas (A04), otras intoxicaciones alimentarias bacterianas (A05), amebiasis (A06), otras enfermedades intestinales debidas a protozoarios (A07), infecciones intestinales debidas a virus (A08) y diarrea y gastroenteritis de presunto origen infeccioso (A09). Se utilizaron datos de acceso abierto de la página WEB de la Dirección General de Información Estratégica en Salud del MSP y BS de todos los Departamentos del país. Se calcularon tasas crudas y ajustadas (por 100.000 habitantes). Se utilizaron Excel y EPI INFO 7.0. **Resultados:** Se registraron 495 muertes, la mayor prevalencia fue en 2019 (tasa ajustada de 1,83) y menor en 2017 (1,36). Los meses con mayor mortalidad fueron enero y julio (9,9%). El 51,72% fueron mujeres (256), 30,91% mayores de 80 años y 28,28% (140) menores de 5 años, edad promedio 50,8, solteros 56,77% (281) y sin ningún tipo de educación 41,82% (207). El 80,61% (399) residía en área urbana. Las mayores tasas se registraron en Boquerón (33,3) y Amambay (15,2). El 95,56% (473) fueron diarreas y gastroenteritis de origen infeccioso. **Discusión:** Se registra alto porcentaje de mortalidad de diarreas de origen infeccioso. Los extremos de la vida constituyen la población más vulnerables.

Palabras clave: gastroenteritis; infecciones; diarrea; Paraguay.

How to cite this article: Mereles-Aranda EF, Tullo-Gómez JE, Duarte-Zolán D, Oliveira Da Silva E, Rivas-Kiese ME, Ramos P. Epidemiological characteristics of mortality from infectious intestinal diseases (diarrhea) Paraguay (2015-2019). Med. clín. soc. 2024;8(1):84-90

INTRODUCTION

Infectious intestinal diseases (diarrhea) are a major cause of morbidity and mortality worldwide. In 2015, it constituted the ninth leading cause of death for all ages; in 2017, in children under five years old, it was the third leading cause of death and has been shown to be the cause of at least 8 million deaths in low- and middle-income countries, with a cost of economic losses of at least \$6 trillion (1-3). In children under 5 years of age, the number of diarrhea episodes per year varies from 3,3 to 9 episodes (4). In addition to mortality, acute or chronic diarrhea affects the development and growth of children, mainly during the first two years of life, correlating with poor school performance and low wages in adulthood (5).

Most diarrheas have infectious etiology, being the main pathogenic agents; viruses (*Norovirus*, *Rotavirus*, *Astrovirus* and enteric *Adenovirus*) bacteria (*Salmonella* spp., *Shigella* spp., *Campylobacter* spp., *Vibrio* spp., *E. coli* diarrhea spp., *Aeromonas* spp., *Yersinia* spp., among others) and, less frequently, parasites (*Cryptosporidium* spp., *Giardia* spp., *Cyclospora* spp., *Microsporidium* spp., *Entamoeba histolytica*, *Balantidium coli*) (5). According to the World Health Organization (WHO), in low-income countries, the most common pathogens in low-income countries are *Rotavirus*, *Escherichia coli*, *Cryptosporidium* spp., and *Shigella* spp. (4).

In general, diarrhea is self-limiting, and although deaths from diarrhea have declined sharply in the past two decades, it is estimated that by 2050, 1.4 million people will die from diarrheal diseases (6), mainly due to the great social and health inequities in low-income countries. Thus, in these countries, the social determinants of health form a complex multifactorial network in which risk factors related to the individual, environment, and animals are intertwined (7). 80% of the risk factors for diarrhea are related to poor environmental conditions such as basic sanitation, hygiene habits, poor quality and low coverage of drinking water, and in children, malnutrition, and dehydration (4,8,9).

Furthermore, in addition to poor environmental conditions, there is a lack of access to and poor quality of health systems. In developed countries, deaths due to diarrhea are infrequent, unlike in underdeveloped countries, where hospital admissions are observed due to complications such as severe dehydration and renal failure, especially in elderly patients. This population may present with diarrhea in up to 20% of cases during the treatment of diabetes mellitus (6). Therefore, in recent years, the use of oral rehydration salts has been

emphasized to prevent dehydration, which is the main cause of mortality in children and the elderly (4). Paraguay ranks the lowest in the Human Development Index (HDI), with a value of 0.728, ranking 103 out of 189 countries and territories. Its ranking is lower than the average of countries with a high HDI and the average of Latin America and the Caribbean (10). Therefore, it presents great social and health inequities, which have repercussions on the prevalence of certain infectious diseases such as diarrhea.

Acute childhood diarrhea constitutes in Paraguay, the fourth cause of mortality in children under 5 years old and according to the General Directorate of Health Surveillance (GDHS) of the Ministry of Public Health and Social Welfare (MSPBS), notifications for diarrhea were increasing from 49,750 cases for the year 2015 to 226,413 for the year 2018. Broadly speaking, except for the year 2015, in the other years, most of the cases were in those older than 5 years (11-15). In other studies, it was observed that 40.1% of the cases corresponded to diarrhea in children under 5 years of age, and 59.8% of the cases were observed in children over 5 years of age (16,17). The objective of this study was to describe mortality due to diarrhea in Paraguay between 2015 and 2019.

METHODS

The study was descriptive, ecological, cross-sectional, and temporally retrospective. The geographical scope was composed of 17 Paraguayan Departments, including Asunción, the capital city. The study period was from 2015 to 2019. Sampling was nonprobabilistic for consecutive cases. The considered variables were sex, marital status, educational level, age, and 17 Paraguayan Departments. The place of residence of the deceased was considered as the origin of the cases. Deaths from non-nationals were excluded from the calculation of the rates.

Deaths due to diarrhea of infectious etiology were considered, which were classified in the ICD-10 as infectious intestinal diseases, including cholera (A01), typhoid and paratyphoid fevers (A02), shigellosis (A03), other bacterial intestinal infections (A04), other bacterial food poisoning not elsewhere classified (A05), amebiasis (A06), other intestinal diseases due to protozoa (A07), intestinal infections due to viruses and other specified organisms (A08), diarrhea, and gastroenteritis of suspected infectious origin (A09) (18). The data source used was the open-access data available on the website of the General Directorate of Strategic Health Information (GDSHI) of the Ministry of Public Health and Social Welfare (MSPyBS) (19).

Crude and adjusted rates (100,000 per capita) were calculated. Excel and EPI INFO 7.0 were used. For statistical analysis of nominal and categorical variables, frequencies were calculated and expressed as percentages. Averages were calculated for the numerical variables. Gross and adjusted rates were also calculated using population data from the National Institute of Statistics (NIS) and the World Health Organization's standardized population standard (20-22). Regarding ethical criteria, as this was a research study with a secondary data source that did not involve human subjects, the protocol of the present study did not require the approval of the Research Ethics

Committee.

RESULTS

From 2015 to 2019, 495 people died of diarrhea of infectious etiology, the main etiology being diarrhea and gastroenteritis of presumed infectious origin (A09), which constituted 95.56% of the deaths. During this period, there were no deaths due to diarrhea caused by cholera (A01), typhoid and paratyphoid fevers (A02), shigellosis (A03), amebiasis (A06), or viruses (A08) (Table 1).

TABLE 1. ETIOLOGY OF DEATHS DUE TO INFECTIOUS INTESTINAL DISEASES (DIARRHEA) ACCORDING TO THE CIE – 10, PARAGUAY (2015-2019) (n=495).

Variables	Frequency	Percentage	IC 95%
Other gastroenteritis and colitis of infectious and unspecified origin (A09)	473	95.56	93.36 – 97.05
Other bacterial intestinal infections (A04)	20	4.04	2.63 – 6.16
Other bacterial foodborne intoxications, not elsewhere classified (A05)	1	0.20	0.04 – 1.14
Other protozoal intestinal diseases(A07)	1	0.20	0.04 - 1.14


The year with the highest number of deaths was 2019, with an adjusted rate of 1.83 per 100,000 people and 23.84% of cases, and the year with the lowest prevalence was 2015, with an adjusted rate of 1.36 per 100,000 people (Table 2). As for the monthly distribution of cases, January and July (9.9%) and October (9.3%) had the highest number of cases

(Figure 1). Regarding the age and sex of the deceased, the highest number of deaths occurred in both the earlier and later stages of life. Thus, in children under 5 years of age, most deaths occurred in males. However, in older adults, from 65 years of age onwards, most deaths occurred in the female sex (Figure 2)

TABLE 2. ANNUAL DISTRIBUTION OF INFECTIOUS INTESTINAL DISEASE (DIARRHEA) CASES, PARAGUAY (2015-2019) (n=495).

Years	Number of cases	Percentage	Cruda rate 100 000 hab.	Adjusted rate 100 000 hab.
2015	81	16.36	1.20	1.36
2016	97	19.60	1.42	1.57
2017	93	18.79	1.34	1.51
2018	106	21.41	1.50	1.67
2019	118	23.84	1.65	1.83

FIGURE 1. MONTHLY PERCENTAGE DISTRIBUTION OF INFECTIOUS INTESTINAL DISEASES (DIARRHEA) CASES PER YEAR, PARAGUAY (2015-2019) (n=495).



The highest number of deaths occurred in both the earlier and later stages of life. Thus, in children under 5 years of age, most deaths occurred in males. However,

in older adults, from 65 years of age onwards, most deaths occurred in the female sex (Figure 2). Regarding the demographic characteristics, 51.72% were women.

In terms of age, the average was 50.8 years. When analyzing the distribution according to age categories, 30.91% of the cases occurred in those over 80 years of age and 28.28% were under 5 years of age (Table 3).

Regarding the area of residence of the deceased, the highest mortality rates occurred in the Departments of Boquerón (33.3 per 100 000 inhabitants) and Amambay (15.2) (Table 4).

FIGURE 2. DISTRIBUTION OF DEATHS FROM INFECTIOUS INTESTINAL DISEASES (DIARRHEA) BY AGE AND SEX, PARAGUAY (2015-2019) (n=495).




TABLE 3. DEMOGRAPHIC CHARACTERISTICS OF PATIENTS WHO DIED FROM INFECTIOUS INTESTINAL DISEASES (DIARRHEA), PARAGUAY (2015–2019) (n=495).

Variables	Frecuency	Percentage	IC 95%
Sex			
Female	256	51.72	47.32 - 56.09
Male	239	48.28	43.91 - 52.68
Age category			
0 – 4	140	28.28	24.49 – 32.41
5 – 9	13	2.63	1.54 – 4.44
10 – 14	4	0.81	0.31 – 2.06
15 – 19	3	0.61	0.21 – 1.77
20 – 24	4	0.81	0.31 – 2.06
25 – 29	2	0.40	0.11 – 1.46
30 – 34	6	1.21	0.56 – 2.62
35 – 39	4	0.81	0.31 – 2.06
40 – 44	10	2.02	1.10 – 3.68
45 – 49	10	2.02	1.10 – 3.68
50 – 54	12	2.42	1.39 – 4.19
55 – 59	15	3.03	1.84 – 4.94
60 – 64	16	3.23	2.00 – 5.19
65 – 69	35	7.07	5.13 – 9.67
70 – 74	30	6.06	4.28 – 8.52
75 – 79	38	7.68	5.64 – 10.36
80 +	153	30.91	27.00 – 35.11
Marital status			
Single	281	56.77	52.37 - 61.06
Married	136	27.47	23.73 - 31.57
Widower	63	12.73	10.08 - 15.95
Separated	3	0.61	0.21 - 1.77
United	3	0.61	0.21 - 1.77
Divorced	2	0.40	0.11 - 1.46
Does not know	7	1.41	0.69 - 2.89
Education level			
None	207	41.82	37.55 - 46.21
Incomplete primary	91	18.38	15.22 - 22.04
Complete primary	84	16.97	13.92 - 20.53
Does not know	64	12.93	10.26 - 16.17
Complete secondary	35	7.07	5.13 - 9.67
Complete University or higher	10	2.02	1.10 - 3.68
Complete Basic school	4	0.81	0.31 - 2.06
Area of residence			
Urban	399	80.61	76.89 - 83.85
Rural	96	19.39	16.15 - 23.11

TABLE 4. MORTALITY RATE DUE TO INFECTIOUS INTESTINAL DISEASES (DIARRHEA) ACCORDING TO THE DEPARTMENT OF RESIDENCE OF THE DECEASED, PARAGUAY (2015-2019) (n=495).

Department	Frequency	Percentage	Rate mortality (100mil hab.)
1. Concepción	28	5.7	11.5
2. San Pedro	17	3.4	4.1
3. Cordillera	21	4.2	7.0
4. Guairá	16	3.2	7.2
5. Caaguazú	40	8.1	7.3
6. Caazapá	12	2.4	6.5
7. Itapuá	26	5.3	4.4
8. Misiones	13	2.6	10.5
9. Paraguari	14	2.8	5.5
10. Alto Paraná	61	12.3	7.7
11. Central	117	23.6	5.6
12. Ñeembucú	7	1.4	7.9
13. Amambay	25	5.1	15.2
14. Canindeyú	10	2.0	4.5
15. Presidente Hayes	18	3.6	14.9
16. Boquerón	21	4.2	33.3
17. Alto Paraguay	2	0.4	11.6
18. Asunción (Capital)	46	9.3	8.8
Foreign	1	0.2	-

DISCUSSION

Infectious diarrhea continues to be one of the main causes of death from diarrhea in Paraguay, and despite efforts, there has been a sustained increase over the years. This could be because notification systems and the quality of death certificates have been improving over time, so that under-registration has been decreasing and the quality of death certificates has improved. Another hypothesis is that this increase is because diarrhea is on the rise due to the persistence of poor conditions related to the environment (hygiene, coverage, and quality of drinking water and sanitary sewage) and to the individual (low or low level of education).

According to data from the Health Services Company of Paraguay (HSCP), drinking water coverage in Paraguay accounts for 80% of the population in urban areas and 33% in rural areas. Drinking water is distributed through a network of pipes that are in very poor condition, causing significant water losses (47% of losses) and leakage inside the network. The HSCP estimates that 300,000 m of pipes must be replaced. The poor quality of these pipes also favors contamination of the water being transported. Another point related to the environment is that there is practically no sanitary sewage system; therefore, there is no wastewater treatment. Asunción, the country's capital, and the metropolitan area have 33% of its population connected to the sanitary sewer system (22-25).

As for factors related to the individual, diarrhea has

moved from the pediatric population to the geriatric population, since it can be observed that the highest prevalence of mortality occurred at both the early and later stages of life, those over 80 years of age, and those under 5 years of age. This is because this population is increasing among those over 80 years of age. According to INE data, this population currently includes 746,445 people aged 60 and over, of whom 48.3% are men and 51.7% are women (26). Even so, it is striking that mortality is due to this pathology since diarrhea, in general, can be cured with good oral rehydration. Further research should be conducted to determine whether this mortality is related to a lack of access to health services.

In this demographic group, acute diarrhea could be due to the prolonged use of broad-spectrum antibiotics or fragility of the immune system (27). Most deaths occurred in women aged > 65 years. This could be because they often live alone, have a low educational level, have little or no work activity, face specific age-related problems, and do not attend medical consultations. It could be because it is women who have a longer life expectancy, as well, with a projected 80.7 years for this sex by 2030 according to the WHO (27,28).

In children under 1 and 5 years of age, Paraguay presented values that placed it above countries such as Haiti (10%), Nicaragua (8%), Guatemala (7%), and Bolivia (6%), with 14.2 and 16.8 deaths per 1,000 live births, respectively, being one of the main causes of death along with pneumonia and influenza (29).

Regarding pathogenic agents, the absence of mortality due to *Salmonella* spp. and *Shigella* spp. is striking, as these agents are the most prevalent in diarrhea of infectious etiology. Evidently, these pathogens are agents of morbidity, but not mortality (30). In Asunción and the metropolitan area, diarrheagenic *Escherichia coli*, Rotavirus, *Campylobacter* spp., *Shigella* spp., and *Salmonella* spp. have been described as the most prevalent etiological agents in children under 5 years of age (31).

The absence of mortality due to cholera is expected, since Paraguay reported only two cases of this pathogen even during the cholera epidemic. Consequently, Paraguay is considered cholera-free (32). However, approximately 1.3 4 million cases of cholera and 21,000–143,000 deaths due to cholera have been reported annually worldwide (33).

Regarding mortality due to food poisoning, this could be considered underreported because of the precarious conditions in which many establishments work in this area and the lack of control by the authorities in charge. Epidemiological surveillance in the last two or three decades has shown an increase in the prevalence of foodborne diseases (FBD), with outbreaks of diseases such as salmonellosis, Enterohemorrhagic *Escherichia coli* infections (EHEC), and hepatitis A. There is little literature on this topic; the only information available reports that between 2015 and 2016, a total of 14 outbreaks of FBD were recorded, with a total of 231 cases, with no deaths reported because of this health event (34,35).

There were numbers concerning the monthly distribution of cases; the months with the highest number of cases were January, July, and October. Bacterial diarrhea is more prevalent in the warm summer months, and since no mortality due to viruses was observed in this five-year period, we assumed that the mortality due to diarrhea in July was also due to bacterial agents. Another point to consider is that Paraguay has high temperatures throughout the year. The month of January coincides with work carried out in Argentina, where there were two peaks during the year. One of the peaks occurred during the warm months, with a predominance of bacterial infections, and the other peak occurred during the cold months, where the prevalence of viral diseases was observed (36).

Regarding the area of residence, the highest mortality rates (33.3 and 15.2) were in the Departments of Boquerón and Amambay, respectively, both geographically distant from the most complex healthcare centers for better care. In most developed

and developing countries, mortality due to diarrhea is no longer a major public health concern. However, in Paraguay, mortality due to diarrhea of infectious etiology continues to be a complex problem that requires an approach based on social determinants of health for its prevention and elimination.

AUTHORS CONTRIBUTIONS

All authors contributed to the study conception and conception, data interpretation, statistical analysis, drafted the article and revised and approved the final version.

REFERENCIAS

1. Kruk ME, Gage AD, Arsenaault C, Jordan K, Leslie HH, Roder-DeWan S, et al. High-quality health systems in the Sustainable Development Goals era: time for a revolution. *The Lancet Global Health*. 2018;6(11):e1196–252. [https://doi.org/10.1016/S2214-109X\(18\)30386-3](https://doi.org/10.1016/S2214-109X(18)30386-3)
2. GBD Diarrhoeal Diseases Collaborators. Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis*. 2017;17(9):909-948. [https://doi.org/10.1016/S1473-3099\(17\)30276-1](https://doi.org/10.1016/S1473-3099(17)30276-1)
3. GBD 2016 Diarrhoeal Disease Collaborators. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of diarrhoea in 195 countries: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Infect Dis*. 2018;18(11):1211-1228. [https://doi.org/10.1016/S1473-3099\(18\)30362-1](https://doi.org/10.1016/S1473-3099(18)30362-1)
4. Argentine Ministry of Health. Plan for Comprehensive Approach to Acute Diarrheal Diseases. Guide for the health team No. 8. January/2011. [URL](https://www.argentina.gob.ar/salud/prevenccion-y-promocion).
5. Riveros M, Ochoa TJ. Enteropathogens of importance in public health. *Rev Peru Med Exp Public Health*. 2015;32(1):135-43. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).
6. Galas M. Antimicrobial resistance: from the laboratory to patient care. 2018. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).
7. Boo YY, Rai K, Cupp MA, Lakhanpaul M, Factor-Litvak P, Parikh P, et al. What are the determinants of childhood infections in India's periurban slums? A case study of eight cities. *PLoS ONE*. 2021;16(10):e0257797. <https://doi.org/10.1371/journal.pone.0257797>
8. Pan American Health Organization (PAHO/WHO). Strengthening of sanitation boards. 2011. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).
9. Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet*. 2016; 388:3027–35. [http://dx.doi.org/10.1016/S0140-6736\(16\)31593-8](http://dx.doi.org/10.1016/S0140-6736(16)31593-8)
10. United Nations Development Program UNDP. The next frontier: human development and the Anthropocene. Information note for countries on the Human Development Report 2020. Human Development Report 2020. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).
11. Weekly Epidemiological Bulletin. General Directorate of Health Surveillance (DGVs). Ministry of Public Health and Social Welfare. Edition No.: 18. May 15, 2015. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).
12. Weekly Epidemiological Bulletin. General Directorate of Health Surveillance (DGVs). Ministry of Public Health and Social Welfare. Edition No. 48. SE 1 (January 1) to SE 52 (December 30). 2018. [URL](https://doi.org/10.1016/S1473-3099(18)30362-1).

13. Weekly Epidemiological Bulletin. General Directorate of Health Surveillance (DGVS). Ministry of Public Health and Social Welfare. Edition No.52. SE 1 a la SE 52 (31-12-2017 al 29-12-2018). 2018. [URL](#).
14. Weekly Epidemiological Bulletin. General Directorate of Health Surveillance (DGVS). Ministry of Public Health and Social Welfare. Edition No.52. SE 1 a la SE 52 (31-12-2017 al 29-12-2018). 2018. [URL](#).
15. Weekly Epidemiological Bulletin. General Directorate of Health Surveillance (DGVS). Ministry of Public Health and Social Welfare. Edition No. SE 1 a la SE 47 (30-12-2018 al 23-11-2019). Pags:23 – 27. 2019. [URL](#).
16. Martínez M, Fariña N, Rodríguez M, Russomando G, Parra GI. Presencia de rotavirus en adultos con diarrea en Asunción, Paraguay. Rev. argent. microbiol. 2005;37(2):99–101. [URL](#).
17. Ministry of Public Health and Social Welfare. Diarrhea can be due to poor hygiene or food poisoning. 2019. [URL](#).
18. Pan American Health Organization International Statistical Classification of Diseases and Health Related Problems. — 10a. revision. Washington, DC: PAHO. 1995. [URL](#).
19. Ministry of Public Health and Social Welfare. General Directorate of Strategic Health Information. Historical Series of Mortality Indicators (INDIMOR).2015-2019. 2019. [URL](#).
20. INE- National Statistics Institute (INE). Projection of the Population by Sex and Age, 2019. 2019. [URL](#).
21. National Statistics Institute. C4. Paraguay. Total population, estimated and projected, by department, sex and age, 2000-2025. 2023. [URL](#).
22. World Health Organization (WHO). Age Standardization of Rates: A new WHO Standard. GPE. Discussion Paper Series N ° 31. 2023. [URL](#).
23. López, M. The older adult population in Paraguay: With reduced access to pensions and submerged in informality. 2021. [URL](#).
24. PAHO. Crecimiento acelerado de la población adulta de 60 años. 2023. [URL](#).
25. STP. Aging Paraguay. [URL](#).
26. United Nations, Department of Economic and Social Affairs, Population Division. World population prospects: the 2017 revision, volume II, demographic profiles. New York: United Nations;2017. [URL](#).
27. World Health Organization. Global strategy and plan of action on aging and health. 69th World Health Assembly, Geneva, April 22, 2016 (A69/17). [URL](#).
28. World Health Organization. Global strategy and plan of action on aging and health. 69th World Health Assembly, Geneva, April 22, 2016 (A69/17). [URL](#).
29. Pan American Health Organization. Health in the Americas, 2017 edition. Summary: regional overview and country profiles. 2017. [URL](#).
30. Díaz SL, Jarquin C, Morales AJ, Morales M, Valenzuela C. Burden of salmonellosis and shigellosis in four departments of Guatemala, 2010. Rev Panam Salud Publica.2015;38(4):326–32. [URL](#).
31. Huber C, Orrego MV, Ortiz F, Álvarez M, Weiler N, Huber C, et al. Prevalencia de patógenos causantes de enfermedad diarreica aguda en el área Metropolitana de Asunción y Central. Rev. salud publica Parag. 2019;9(2):41–5. <https://doi.org/10.18004/rspp.2019.diciembre.41-45>
32. Ramirez, ME. Brief tour of the great pandemics and epidemics that hit the world and Paraguay.2020. [URL](#).
33. WHO. Cholera. 2021. [URL](#).
34. Sotelo AB, Martínez C, Sánchez S, Sotelo AB, Martínez C, Sánchez S. Características epidemiológicas y clínicas de los brotes de Enfermedades Transmitidas por Alimentos. Paraguay 2015 - 2016. Rev. salud publica Parag. 2019;9(1):33–40. <https://doi.org/10.18004/rspp.2019.junio.33-40>
35. PAHO. MSP and BS. Foodborne Disease Surveillance Manual. 2017. [URL](#).
36. Degiuseppe JI. Trend of hospital discharges for intestinal infectious diseases in children in Argentina, 2005-2013. Arch Argent Pediatr 2017;115(4):350-356. <http://dx.doi.org/10.5546/aap.2017.350>