


The effect of Orem self-care model on the improvement of symptoms and quality of life in patients with diabetes: A scoping review

Mohammadamin Jandaghian-Bidgoli¹ 
<https://orcid.org/0000-0003-0195-9099>

Sheida Jamalnia^{2,3} 
<https://orcid.org/0000-0001-5923-5822>

Marzieh Pashmforosh⁴ 
<https://orcid.org/0000-0002-6697-3396>

Negin Shaterian⁵ 
<https://orcid.org/0000-0003-1827-941X>

Pouriya Darabiyan⁶ 
<https://orcid.org/0000-0002-9529-1730>

Alireza Rafi⁷ 
<https://orcid.org/0000-0001-5442-2741>

The effect of Orem self-care model on the improvement of symptoms and quality of life in patients with diabetes: A scoping review

Abstract

Objective: to evaluate the association of Orem self-care model improvement of symptoms and quality of life in patients with diabetes. **Methods.** A scoping review was carried on bibliographic databases: PubMed-Medline, Scopus, SID and Magiran. The inclusion criteria encompassed studies examining the impact of the Orem self-care model on diabetic patients. Studies considered

- 1 Student Research Committee, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: mohammadaminjandaghian@sbmu.ac.ir
- 2 Department of Nursing and Midwifery, Kazeroun Branch, Islamic Azad University, Kazeroun, Iran.
- 3 Ph.D student of e-Learning, virtual School, Shiraz University of Medical Sciences, Shiraz, Iran. Email: shjamalnia1988@gmail.com
- 4 Behbahan Faculty of Medical Sciences, Behbahan, Iran. Email: marzie_pf@yahoo.com
- 5 Student Research Committee, Kashan University of Medical Sciences, Kashan, Iran. Email: negin_shaterian@yahoo.com
- 6 Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: Pouriya.d.k.76@gmail.com
- 7 M.Sc of Nursing, Behbahan Faculty of Medical Sciences, Behbahan, Iran. Email: alirezaraf72@gmail.com. (Corresponding author)

Conflicts of interest: No

Received: July 16, 2023

Approved: February 2, 2024.

How to cite this article: Jandaghian-Bidgoli M, Jamalnia S, Pashmforosh M, Shaterian N, Darabiyan P, Rafi A. The effect of Orem self-care model on the improvement of symptoms and quality of life in patients with diabetes: A scoping review. *Investigación y Educa. Enferm.* 2023; 42(1):e08.

DOI: <https://doi.org/10.17533/udea.iee.v42n1e08>



<https://creativecommons.org/licenses/by-nc-sa/4.0>



Review



UNIVERSIDAD
DE ANTIOQUIA
1803

Investigación y Educación en

Enfermería

Vol. 42 No 1, January - April 2024
ISSNp: 0120-5307 • ISSNe: 2216-0280

for inclusion needed to have full-text availability and be written in either English or Persian, with key words including “Models”, “Nursing”, “Quality of Life”, and “Diabetes Mellitus”. CONSORT checklist and STROBE statement were selected for quality assessment. **Results.** A total of 9 studies were included, all using quantitative methodology and focusing on adults or older adults. The majority of articles focused on quality of life and diabetic symptoms. 8 studies showed positive outcomes after implementation of the model. The findings indicate that this model led to an enhanced level of self-efficacy, improved quality of life, and better self-care practices among diabetic patients. **Conclusion.** Orem self-care model can reduce the diabetic symptoms and improve the quality of life, self-efficacy and self-care in these patients.

Descriptors: Diabetes Mellitus; models, nursing; self-care; quality of life.

Efecto del modelo de autocuidado de Orem en el mejoramiento de los síntomas y en la calidad de vida de pacientes con diabetes: Una revisión de alcance

Resumen

Objetivo. Evaluar la asociación del modelo de autocuidado de Orem en el mejoramiento de los síntomas y en la calidad de vida en pacientes con diabetes.

Método. Se realizó una revisión de alcance empleando las bases bibliográficas PubMed-Medline, Scopus, SID y Magiran. Los criterios de inclusión abarcaron estudios que examinaran el impacto del modelo de autocuidado de Orem en pacientes diabéticos. Los estudios considerados para su inclusión debían tener disponibilidad de texto completo y estar escritos en inglés o persa, con palabras clave como: “Models”, “Nursing”, “Quality of Life” y “Diabetes Mellitus”. Se utilizaron para la evaluación de la calidad de los estudios la lista de comprobación CONSORT y la declaración STROBE. **Resultados.** Se incluyeron un total de 9 estudios, todos ellos con metodología cuantitativa y centrados en adultos y en ancianos. La mayoría de los artículos se estudiaron la calidad de vida y los síntomas diabéticos. 8 estudios mostraron resultados positivos tras la aplicación del modelo de Orem. Los hallazgos

indicam que este modelo condujo a un mayor nivel de autoeficacia, mejor calidad de vida y mejores prácticas de autocuidado entre los pacientes diabéticos. **Conclusión.** El modelo de autocuidado de Orem puede ayudar a disminuir los síntomas diabéticos y mejorar la calidad de vida, la autoeficacia y el autocuidado en estos pacientes.

Descriptores: Diabetes Mellitus; modelos de enfermería; calidad de vida; autocuidado.

Efeito do modelo de autocuidado de Orem na melhora dos sintomas e na qualidade de vida de pacientes com diabetes: uma revisão de escopo

Resumen

Objetivo. Avaliar a associação do modelo de autocuidado de Orem na melhora dos sintomas e na qualidade de vida de pacientes com diabetes. **Métodos.** Foi realizada uma revisão de escopo usando os bancos de dados PubMed-Medline, Scopus, SID e Magiran. Os critérios de inclusão incluíram estudos que examinaram o impacto do modelo de autocuidado de Orem em pacientes diabéticos. Os estudos considerados para inclusão tinham que estar disponíveis em texto completo e escritos em inglês ou persa, com palavras-chave como: “Models”, “Nursing”, “Quality of Life” e “Diabetes Mellitus”. A lista de verificação CONSORT e a declaração STROBE foram usadas para avaliar a qualidade dos estudos. **Resultados.** Foram incluídos 9 estudos, todos com metodologia quantitativa e com foco em adultos e idosos. A maioria dos artigos estudou a qualidade de vida e os sintomas diabéticos. Oito estudos mostraram resultados positivos após a aplicação do modelo de Orem. Os achados indicam que esse modelo levou a um nível mais alto de autoeficácia, melhor qualidade de vida e melhores práticas de autocuidado entre os pacientes diabéticos. **Conclusão.** O modelo de autocuidado de Orem pode ajudar a diminuir os sintomas da diabetes e melhorar a qualidade de vida, a autoeficácia e o autocuidado desses pacientes.

Descritores: Diabetes Mellitus; modelos de enfermagem; qualidade de vida; autocuidado.

Introduction

Diabetes originates from impairment in the metabolism of proteins, carbohydrates and lipids.⁽¹⁾ It is a chronic metabolic disorder that is defined and recognized by the increase in blood sugar caused by defects in the secretion or action of insulin or both. This disease leads to many complications such as limb amputation, cardiovascular problems, kidney diseases, blindness, and long-term disabilities.⁽²⁾ Diabetes is one of the most common non-communicable diseases that has the highest prevalence rate among metabolic diseases. More than 90% of diabetic patients suffer from type 2 diabetes. This disease is associated with short-term and long-term complications that are irreversible in many cases.⁽³⁾ Complications and deaths caused by diabetes are among health problems all over the world. The increase in the number of patients with type 2 diabetes indicates a global epidemic.⁽⁴⁾ In addition to direct complications, diabetes also has many costs that affect the lives of the patients. In 2017, 327 billion dollars was the total cost of diagnosed diabetes.⁽⁵⁾

The prevalence of type 1 and type 2 diabetes is increasing worldwide, but the rate of type 2 diabetes is higher than that of type 1.⁽⁶⁾ Diabetes is a chronic condition that imposes major threats to the mankind and affected 435 million individuals in 2017.⁽⁷⁾ It will involve 552 and 624 million patients by 2030 and 2040, respectively.^(8,9) According to the International Diabetes Federation (IDF), Iran ranks third among Middle Eastern countries in terms of diabetes (5.4 million cases). It is expected that the rate of diabetes will reach 9.2 million Iranians by 2030.⁽¹⁰⁾ Aging population will be another effective factor in increasing the prevalence of diabetes in the future.⁽¹¹⁾ The increase in the number of diabetic patients has faced the care provider organizations with increasing financial problems and a decrease in the number of care providers.⁽¹²⁾ 15-20% of hospital beds are occupied by patients with diabetes and it is difficult to determine which of these patients require more medical attention.⁽¹³⁾ Regardless of age, country and economic conditions, diabetes is a major health challenge and has a global impact, so that the prevalence of diabetes is reaching an alarming level.⁽¹⁴⁾ Therefore, the prevention and control of this disease is considered one of the health challenges, so that one of the goals of the Ministry of Health, Treatment and Medical Training of Iran is to stop the spread and growth of diabetes until 2025.⁽¹⁵⁾

One of the most important clinical goals of nursing care is to reduce the severity of disease symptoms and the stress caused by it in order to improve the quality of life in these patients. Without patient participation, favorable outcomes cannot be considered. Quality of life is a crucial aspect of healthcare, particularly in the context of nursing care. The primary clinical objective for nurses is to alleviate the severity of disease symptoms and mitigate the associated stress experienced by patients. By doing so, the overarching goal is to enhance the overall quality of life for individuals undergoing medical

treatment.⁽¹⁶⁾ Self-care behaviors are one of the most important components which requires special attention. These behaviors in diabetic patients includes appropriate glycemic control, following a healthy diet, taking medicines regularly, monitoring blood sugar, taking care of feet and doing sports activities.⁽¹⁷⁾ Self-care behaviors are naturally multidimensional and include all the activities focused on health maintenance, disease prevention and treatment. These kinds of behaviors are done purposefully and with the consent of the patient.⁽¹⁸⁾ Improving the self-care behaviors is the first step toward helping the diabetic patients. Normally, self-care practices can reduce the costs and prevent acute or chronic complications of diabetes.^(19,20) Appropriate behaviors can lead to a decreased level of cardiovascular risks among diabetic patients.⁽²¹⁾ Since self-care activities are effective in glycemic control and other outcomes in diabetic patients, it can be claimed that self-care behaviors can be an underlying reason for mortality among them.^(22, 23)

One of the models that can contribute to the patient's self-care is Orem self-care model.⁽²⁴⁾ According to Orem, humans have the ability to take care of themselves, and if this ability is disturbed, nurses help them to regain this ability.⁽²⁵⁾ Understanding the impact of the Orem Self-Care Model on symptom alleviation and quality of life is crucial for healthcare professionals and policymakers. Positive findings could lead to the development of targeted interventions and strategies that enhance patient outcomes, potentially improving the overall quality of care for individuals with diabetes. The need for research on factors affecting self-care based on the needs of patients to increase their self-care ability seems necessary. Hence, this study aimed to evaluate the association of Orem self-care model improvement of symptoms and quality of life in patients with diabetes.

Methods

This scoping review adheres to PRISMA guidelines. The review was registered on PROSPERO

(ID:CRD42022367809). (ID:CRD42022367809) Also, this research has been approved by the ethics committee of Behbahan Faculty of Medical Sciences (code of ethics: IR.BHN.REC.1402.002)

Search strategy. The following databases were searched: PubMed, Scopus, SID, Magiran. Articles published between 2011 and October 31, 2022 were included in the search strategy. The search included following terms: 1- Orem Self-Care Model (Text Word) OR Model, Orem Self-Care (Mesh Term) OR Self-Care Model, Orem (Mesh Term) OR Orem Self Care Model (Mesh Term); 2- Diabetes Mellitus (Text Word) OR Diabetes Mellitus, Noninsulin-Dependent (Mesh Term) OR Diabetes Mellitus, Ketosis-Resistant (Mesh Term) OR Stable Diabetes Mellitus (Mesh Term) OR Type 2 Diabetes Mellitus (Mesh Term) OR Noninsulin Dependent Diabetes Mellitus (Mesh Term) OR NIDDM (Mesh Term) OR Adult-Onset Diabetes Mellitus (Mesh Term) OR IDDM (Mesh Term) OR Type 1 Diabetes OR Autoimmune Diabetes (Mesh Term) OR Juvenile-Onset Diabetes (Mesh Term) OR Sudden-Onset Diabetes Mellitus; and 3- 1 AND 2. In order to minimize the number of missing data, Google Scholar was also searched. The search strategy was completed by snowball search (e.g., existing systematic reviews and included articles) of included studies.

Inclusion/Exclusion Criteria. The inclusion criteria were as following: Studies published in English or Persian with full text available, participants aged 10-65 years, participants with the diagnosis of diabetes mellitus at any stage of the disease. Included interventions were those that aimed to implement Orem model. Moreover, the intervention can be emanated from either inpatient or outpatient setting. Additionally, randomization may reduce the risk of bias, it may not be fully possible in health care. Hence, the studies with nonrandomized or one-group study designs were included.

The exclusion criteria were as following: interventions of models other than Orem model; intervention specifically aimed to treat homeless

patients, participants with physical or mental handicap and patients involved in forensic process. Gray literature which did not publish as articles, such as posters, organizational projects, class presentations, health messages and other similar literature did not meet the eligibility criteria.

Quality Assessment. For the assessment of observational studies, STROBE statement (Strengthening the Reporting of Observational studies in Epidemiology) was used. The statement consists of 22 items.⁽²⁶⁾ In addition, CONSORT checklist was used for the assessment of interventional studies.⁽²⁷⁾

Data Extraction. In order to find eligible studies, titles and abstracts from databases and additional sources were assessed by two authors independently. Then, the retrieved full texts were checked by the same two authors. Any disagreement was resolved by consultation with the third author. Unavailable data was requested from study authors.

Two authors were responsible for data extraction of included studies. The following items were extracted: 1- General information (Author, publication year, reference, study type, mean age, sample size); 2- Participants; 3- Measurement tool; 4- Intervention period, 4- Control; 5- Outcome; 6- Main finding or key points.




Figure 1. PRISMA Flowchart of selected studies

Results

The initial search yielded 259 records on PubMed, 226 on Scopus, 215 on SID and 189 on Magiran. After removal of duplicates, the final records were 625. By reviewing the titles, the eligible articles reduced to 321. The selection by abstract brought the total to 74. The main purpose of the study was to select the studies have examined the special association between Orem self-care model and diabetic patients. This led to a significant reduction of eligible articles. Eventually, 9 studies met the inclusion criteria and were considered. Figure 1 shows the process of study selection.

Two studies were from PubMed, whilst 6 were extracted from SID and Magiran. Additionally, one study gained from snowball method. Almost, all of the studies were interventional. There were 597 participants in this systematic review.

Within the selected studies, there were some items investigated: 3 articles focus on diabetes-related psychological symptoms,^(28,29) 3 articles focus on self-efficacy,⁽³⁰⁾ 2 articles focus on quality of life,⁽³¹⁻³³⁾ 1 articles focus on self-care.⁽³⁴⁻³⁶⁾ 8 out of 9 articles showed a positive statistically significant correlation between Orem self-care model and positive outcomes in diabetic patients.

The details and main findings are given in Table 1. For instance, it was found that implementing Orem's self-care model can lead to a decreased level of diabetic neuropathy.⁽²⁸⁾ As well, the study showed that the group receiving Orem's model has more appropriate level of self-care.⁽²⁹⁾ Similarly, as previously mentioned, other factors such as quality of life⁽³¹⁻³³⁾ and self-efficacy⁽³⁰⁾ had improved in the group receiving Orem's self-care model.

An overview of quality assessment of studies using STROBE statement and CONSORT checklist and is presented in Tables 2. For the quality assessment of observational studies, the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) was used. STROBE consists of a checklist with 22 items. The total scores lower than 17 were indicative of low methodological quality.⁽³⁷⁾ On the other hand, experimental and quasi-experimental studies were assessed by the CONSORT checklist. This checklist includes 25 items (38). Two authors were responsible for methodological quality of included studies. The third author resolved any disagreement between the reviewers. Seemingly, the included studies possessed acceptable quality.

Table 1. CONSORT and STROBE checklist scores of selected studies

Author and year of publication	Reference	Type of study	Score
Ahrary <i>et al.</i> (2020)	29	Randomized controlled trial	20/24*
Hemmati Maslarpak <i>et al.</i> (2017)	30	Controlled trial	21/24*
Mansouri <i>et al.</i> (2017)	31	Controlled trial	22/24*
Baraz <i>et al.</i> (2017)	32	Controlled trial	22/24*
Borji <i>et al.</i> (2017)	33	Controlled trial	21/24*
Shahbaz <i>et al.</i> (2016)	35	Controlled trial	20/24*
Ganjlo <i>et al.</i> (2015)	34	Randomized controlled trial	21/24*
Ghafourifard <i>et al.</i> (2015)	36	Controlled trial	22/24*
Khosravan <i>et al.</i> (2015)	37	Descriptive	20/22†

*: CONSORT score; †: STROBE score

Discussion

The results indicated Orem self-care model can lead to an improved level of self-efficacy, quality of life and self-care among the diabetic patients. Also, the diabetic symptoms diminished. In this way, similar studies have showed that self-care behaviors can lead to the improvement of quality of life and reduction of costs.⁽³⁹⁾ It was proved that Orem self-care model is effective in changing the dietary habits among the participants,⁽³⁵⁾ as other studies have shown the efficacy of educational programs for improving the dietary behaviors and controlling the blood sugar are effective.⁽⁴⁰⁾ However, the seasonal preferences need to be taken into account. For instance, in autumn, the desire for protein and fat increases, whilst patients in summer prefer to consume more carbohydrate.⁽⁴¹⁾ Even, religious beliefs, culture and lifestyle can affect dietary habits. In other words, dietary habits are individualized.⁽⁴²⁾ Even though, culture can alter the perception of patients regarding healthcare providers.⁽⁴³⁾

Some of the included studies were conducted on patients with diabetic foot ulcer and showed promising results.^(29,34) Other similar programs conducted in Iran were indicative of effectiveness of self-care training programs among diabetic patients as they could be helpful in wound healing process.⁽⁴⁴⁾ Similarly, studies in other countries highlighted the pivotal role of educating patients for the self-care.⁽⁴⁵⁾ It is proved that controlling the diabetic foot ulcer mainly depends on the patient.⁽⁴⁶⁾ However, the factors such as marital status, educational level, gender, age and occupation play important role in the level of self-care.⁽⁴⁷⁾ For instance, it is claimed that rate of self-care is higher in females compared with males.⁽⁴⁸⁾ Likewise, single patients pay less attention to their self-care, especially regarding diabetic foot ulcer.⁽⁴⁸⁾ It is because stronger emotional and social bonds exist among married patients.⁽⁴⁹⁾

The study participants were middle-aged and older adults. In this way, a study on older adults

with diabetes showed health-related quality of life was moderate.⁽⁵⁰⁾ Though, married older adults had higher scores of quality of life.⁽⁵¹⁾ Education was another multi-dimensional factor which can lead to higher contribution in the areas such economy, society and politics. Educational level directly affects the quality of life in the diabetic patients.⁽⁵²⁾ On the other hand, illiterate individuals are less familiar with coping strategies. Also, these individuals are in an inappropriate economic situation, which can cause more stressful situation and lower level of quality of life.⁽⁵³⁾

The results indicated that Orem self-care model in female patients with diabetes suffering from neuropathy can be effective in reeducation of symptoms and severity and it led in improvement of knowledge, attitude and other related skills.^(28,36) However, their performance and self-care abilities was reported moderate.⁽³⁶⁾ Other studies conducted in Iran have also shown that training can be effective in communication skills and self-control in patients with diabetic neuropathy.⁽⁵⁴⁾ In addition, a study in USA proved that education is effective in regulation of cognitive and emotional states.⁽⁵⁵⁾

The psychological effect of self-care programs is also undeniable. For instance, a study showed that self-management were effective in improvement of mood condition in diabetic patients.⁽⁵⁶⁾ However, many programs do not consider education in the area of psychological skills, as problems such as depression are common in diabetic patients.⁽⁵⁷⁾ A study in Iran showed psychological training can alter the level of stress, anxiety and depression among diabetic patients.⁽⁵⁸⁾ Additionally, existential need in patients encourages them to seek meaning in life and this can help to increase their well-being.⁽⁵⁹⁾ In explaining the role of the spiritual dimension of patients, it can be acknowledged that patients who receive more services have more ability to deal with psychological pressure and psychological trauma.⁽⁶⁰⁾ For the success in implementing such programs, the imperative role of economy should not be neglected. Economic

well-being and availability of financial services can be effective in medical adherence of patients.⁽⁶¹⁾ Naturally, unemployed individuals or those with a low-income profession are more vulnerable to experiencing adverse effects of the disorders.⁽⁶²⁾ Additionally, comorbidities play an important part in lack of self-care behaviors in diabetic patients.⁽⁶³⁾ Obstacles such as physical limitations, polypharmacy, adverse symptoms and lack of social and emotional support exist that can have negative impact on self-control and self-care behaviors.⁽⁶⁴⁾ However, it was claimed that increasing comorbidities result in poor diabetes self-care.⁽⁶⁵⁾

One of the findings was focused on gestational diabetes, which indicated Orem self-care model can be effective for the patients.⁽³⁰⁾ Similarly, other studies also proved that self-care training can bring about positive outcome both in Iran and other countries.^(66,67) However, one study claimed that education cannot alter the scores related to blood sugar.⁽⁶⁸⁾ Specially, after COVID-19 pandemic, the psychological problems have increased, which it may be reduced in gestational diabetes by self-care training.^(66,69) Notably, lack of knowledge regarding health is a risk factor for poor self-management in women suffering from gestational diabetes.⁽⁷⁰⁾ Likewise, It is believed that lack of social support is another important risk factor for women and has adverse effects on gestational outcomes.⁽⁷¹⁾ One of the important steps toward coping behaviors in gestational

diabetes is acceptance. Emotional stability is a factor which may lead to the acceptance.⁽⁷²⁾

In conclusion, looking at the relation found between Orem self-care model and diabetic patients' outcome, it can be stated the model can be effective for the diabetic patients. Positive and supportive implementation of the model can improve patients' quality of life, self-efficacy, and self-care behaviors and reduce diabetic symptoms. The studies analyzed in this review have revealed that Orem self-care has a significant association with positive outcomes among diabetic patients. This shows that Orem self-care model can induce transformational alterations in self-care and quality of life in diabetic patients. However, future studies need to focus on other outcomes related to diabetes as it is a multi-dimensional disease. Specially, psychological outcomes need to be considered.

Limitations. Despite the efforts to present a perfect systematic review, there were some shortcomings. In practice, the studies were included if were published in English or Persian owing to inclusion criteria. Also, the full text of one article was not available. Additionally, there were various tools for the measurement of outcomes. Although the search strategy was designed precisely and snowball search of included studies was considered, there may be some missing data. Absolutely, there is need for conducting more studies to complement the findings of current study.

Table 1. Overview of Included Studies

Author, Year, Ref.	Sample Size/ Gender	Study type	Partici- pants	Mean age	Measurement tool	Inter- ven- tion Period	Control	Outcome	Key points/ Main findings
Ahrary <i>et al.</i> (2020) ⁽²⁸⁾	n=120	RCT	Women with type 2 diabetes and diabetic peripheral neuropathy	54.62 ± 6.93	Diabetes Self-Care Activities and Diabetic Peripheral Neuropathy Self-Care, Michigan Neuropathy Screening Instrument, Toronto Clinical Scoring System, Peripheral Neuropathy Self-Care Requisites	Four weeks	Routine care program in the diabetes clinic	Symptoms and severity of diabetic neuropathy, fasting blood sugar and glycosylated hemoglobin	Decreased level of diabetic neuropathy symptoms and severity among participants (P=0.001), glycosylated hemoglobin (P=0.004), fasting blood sugar (P=0.001)
Hemmati Maslakpak <i>et al.</i> (2017) ⁽²⁹⁾	n=50 F:27, M:23	Quasi-experimental	Patients with diabetic foot ulcers (Either Type 1 diabetes or Type 2 diabetes)	54.13 ± 7.64	Self-care status, Saint Elian Wound Score System (SEWSS), Orem assessment form	12 weeks	Usual care	Ischemia, infection, edema, neuropathy, location, topographic aspects, number of affected zones, depth, wound healing phase and self-care	Better scores for self-care, number of affected zones, ischemia, infection and wound healing phase in the intervention group compared with the intervention (P<0.05)
Mansouri <i>et al.</i> (2017) ⁽³⁰⁾	n=42	Quasi-experimental	Women with gestational diabetes mellitus (n=42)	49	Diabetes Self-efficacy Scale (CIDS)	NR	N/A (pre and post type)	Self-efficacy	Higher level of self-efficacy in the intervention group (P=0.01) Better post-intervention score compared with the pre-intervention (P>0.001)

Table 1. Overview of Included Studies. (Cont.)

Author, Year, Ref.	Sample Size/ Gender	Study type	Partici- pants	Mean age	Measurement tool	Inter- ven- tion Period	Control	Outcome	Key points/ Main findings
Baraz <i>et al.</i> (2017) ⁽³¹⁾	n=30 F:15, M:15	Quasi-experi- mental	Patients with Type 2 Diabe- tes	46.69 ± 7.56	Iranian Short Form Health Survey, Quality of Life Questionnaire	110 hours (two ses- sions)	N/A (pre and post type)	Quality of life	Improvement in some of the quality of life compo- nents such as General health (P = 0.027), Physical role (P < 0.001), Physical functioning (P = 0.027), Social functioning (P = 0.029) and Body pain (P = 0.020) No significant difference between the groups regarding emotional role, strength and vital energy, and general health perception
Borji <i>et al.</i> (2017) ⁽³²⁾	n=80 F:36, M:44	Quasi-experi- mental	Patients with Type 2 Diabe- tes	43.80 ± 11.93	The 36-Item Short Form Health Survey (SF-36), Orem assessment form	360 to 540 minutes (6 ses- sions)	No special care	Quality of Life	Significant improvement of quality of life after intervention (P<0.001) No significant improvement in the control group (P>0.05)
Shahbaz <i>et al.</i> (2016) ⁽³⁴⁾	n=60 F:27, M:23	Quasi-experi- mental	Patients with dia- betic foot ulcer	54.57 ± 6.04	Orem assessment form	210- 255 minu- tes (5 ses- sion)	NR	Self-care	Significant diffe- rence between the groups after intervention (P<0.001)

Table 1. Overview of Included Studies. (Cont.)

Author, Year, Ref.	Sample Size/ Gender	Study type	Partici- pants	Mean age	Measurement tool	Inter- ven- tion Period	Control	Outcome	Key points/ Main findings
Ganjloo <i>et al.</i> (2015) ⁽³³⁾	n=75 F:38, M:37	RCT	Patients with diabetes type 2	35-65	Audit of Diabetes Dependent Quality of Life (ADDQoL-19) and Self-care inventory (SCI)	4 weeks	Under the current education of diabetes center	Quality of Life	Improved level of quality of life score after evaluation of social, psycho-spiritual and social dimensions (P>0/001) Significant difference in quality of life score between intervention and comparison group (P>0/001)
Ghafou- rifard <i>et al.</i> (2015) ⁽³⁵⁾	n=20 F:3, M:17	quasi- experi- mental	Hospi- talized diabetic patients	42.64 ± 20.15	Summary of Diabetes Self-care Activities (SDSCA)	NR	N/A (pre and post type)	Self-care	Improved level of score regarding diet (P=0.04), physical activity (P=0.000), blood sugar levels (P=0.003), medication therapy (P=0.001) and diabetic foot care (P=0.01) Improved level of self-care among inter- vention group participants (P>0/05)
Khos- ravan <i>et al.</i> (2015) ⁽³⁶⁾	n=120	Cross- section- al	Women with diabetic periph- eral neuro- pathy	53.50 ± 6.82	Michigan Neuropathy Screening Instru- ment and Toronto Clinical Neuro- pathy Score, Summary of Diabetes Self-care Activities (SDSCA), Toronto Clinical Scoring System	N/A	N/A	Self- care in relation with knowledge, attitude and performan- ce	Weak ability of self-care in the desired outco- mes after based on the Orem self-care model

RCT: Randomized controlled Trial; NR: Not reported; N/A: Not Applicable ; F: Female ; M:Male

References

1. Kelly SD, Neary SL. Ominous octet and other scary diabetes stories: the overview of pathophysiology of type 2 diabetes mellitus. *Physician Assistant Clinics*. 2020; 5(2):121-33. <https://doi.org/10.1016/j.cpha.2019.11.002>
2. 2Georga EI, Protopappas VC, Bellos CV, Fotiadis DI. Wearable systems and mobile applications for diabetes disease management. *Health and Technology*. 2014; 4(2):101-12. <https://doi.org/10.1007/s12553-014-0082-y>
3. Organization WH. Global report on diabetes: World Health Organization; 2016.
4. Organization WH. Plan of action for the prevention and control of noncommunicable diseases in the Eastern Mediterranean Region. 2011.
5. Economic Costs of Diabetes in the U.S. in 2017. *Diabetes Care*. 2018; 41(5):917-28. <https://doi.org/10.2337/dci18-0007>
6. Vala M, Razmandeh R, Rambol K, Nasli Esfahani E, Ghodsi Ghasemabadi R. Mindfulness-based stress reduction group training on depression, anxiety, stress, self-confidence and hemoglobin A1C in young women with type 2 diabetes. *Iranian Journal of Endocrinology and Metabolism*. 2016; 17(5):382-90.
7. Karimy M, Koohestani HR, Araban M. The association between attitude, self-efficacy, and social support and adherence to diabetes self-care behavior. *Diabetology & metabolic syndrome*. 2018; 10(1):1-6. <https://doi.org/10.1186/s13098-018-0386-6>
8. Zaleska-Żmijewska A, Wawrzyniak ZM, Dąbrowska A, Szaflik JP. Adaptive optics (rtx1) high-resolution imaging of photoreceptors and retinal arteries in patients with diabetic retinopathy. *Journal of Diabetes Research*. 2019: 9548324. <https://doi.org/10.1155/2019/9548324>
9. Cho NH, Shaw J, Karuranga S, Huang Y, da Rocha Fernandes J, Ohlrogge A, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes research and clinical practice*. 2018; 138:271-81. <https://doi.org/10.1016/j.diabres.2018.02.023>
10. Moazzam-Jazi M, Najd Hassan Bonab L, Zahedi AS, Daneshpour MS. High genetic burden of type 2 diabetes can promote the high prevalence of disease: A longitudinal cohort study in Iran. *Scientific Reports*. 2020; 10(1):1-11.
11. Ahmadi A, Hasanzadeh J, Rahimi M, Lashgari L. Factors affecting the quality of life in patients with type 2 diabetes Chahar Mahal Bakhtiari. *Journal of North Khorasan University of Medical Sciences*. 2011;3(1):7-13.
12. Peeters JM, Mistiaen P, Francke AL. Costs and financial benefits of video communication compared to usual care at home: a systematic review. *Journal of Telemedicine and Telecare*. 2011; 17(8):403-11. <https://doi.org/10.1258/jtt.2011.110405>
13. Nirantharakumar K, Chen YF, Marshall T, Webber J, Coleman JJ. Clinical decision support systems in the care of inpatients with diabetes in non-critical care setting; systematic review. *Diabetic medicine*. 2012; 29(6):698-708. <https://doi.org/10.1111/j.1464-5491.2011.03540.x>
14. Istepanian RSH, Al-Anzi TM. m-Health interventions for diabetes remote monitoring and self management: clinical and compliance issues. *Mhealth*. 2018; 4:4. [10.21037/mhealth.2018.01.02](https://doi.org/10.21037/mhealth.2018.01.02)
15. Ahmadi Z, Sadeghi T, Loripoor M, Khademi Z. Comparative assessment the effect of self-care behavior education by health care provider and peer on HbA1c level in diabetic patients. *Iranian Journal of Endocrinology and Metabolism*. 2017; 19(3):144-50.
16. Rahimi A, Ahmadi F, Gholiaf M. Effects of applying continuous care model (CCM) on stress, anxiety and depression in hemodialysis patients. *Research in Medicine*. 2006; 30(4):361-9.
17. Shibayama T, Tanha S, Abe Y, Haginoya H, Rajab A, Hidaka K. The role of illness schemata in self-care behaviors and glycemic control among patients with type 2 diabetes in Iran. *Primary Care Diabetes*. 2019; 13(5):474-80. <https://doi.org/10.1016/j.pcd.2019.03.002>
18. Schrauben SJ, Cavanaugh KL, Fagerlin A, Ikizler TA, Ricardo AC, Eneanya ND, et al. The relationship of disease-specific knowledge and health literacy with the uptake of self-care behaviors in CKD. *Kidney International Reports*. 2020; 5(1):48-57. <https://doi.org/10.1016/j.ekir.2019.10.004>
19. Srinath K, Basavegowda M, Tharuni NS. Diabetic self care practices in rural Mysuru, southern Karnataka, India—a need for diabetes self management educational (DSME) program. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2017;11:S181-S6. <https://doi.org/10.1016/j.dsx.2016.12.029>

20. Karota E, Purba JM, Simamora RH, Siregar CT. Use of King's theory to improve diabetics self-care behavior. *Enfermería Clínica*. 2020; 30:95-9. <https://doi.org/10.1016/j.enfcli.2019.12.035>
21. He X, Li J, Wang B, Yao Q, Li L, Song R, et al. Diabetes self-management education reduces risk of all-cause mortality in type 2 diabetes patients: a systematic review and meta-analysis. *Endocrine*. 2017; 55(3):712-31. <https://doi.org/10.1007/s12020-016-1168-2>
22. Chai S, Yao B, Xu L, Wang D, Sun J, Yuan N, et al. The effect of diabetes self-management education on psychological status and blood glucose in newly diagnosed patients with diabetes type 2. *Patient education and counseling*. 2018; 101(8):1427-32. <https://doi.org/10.1016/j.pec.2018.03.020>
23. Gæde P, Lund-Andersen H, Parving H-H, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *New England Journal of Medicine*. 2008; 358(6):580-91. <https://doi.org/10.1056/nejmoa0706245>
24. Otte JL, Carpenter JS. Theories, models, and frameworks related to sleep-wake disturbances in the context of cancer. *Cancer nursing*. 2009; 32(2):90. [10.1097/01.NCC.0000339261.06442.7d](https://doi.org/10.1097/01.NCC.0000339261.06442.7d)
25. Oshvandi K, Keshmiri K, Salavati M, Emkanjoo Z, Musavi S. Effectiveness of education based on Orem's self-care model in self-care activity of patients with implantable cardioverter defibrillators. *Hayat*. 2014; 19(3):47-55.
26. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *International Journal of Surgery*. 2014; 12(12):1495-9. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X)
27. Moradi M-T, Asadi-Samani M, Mobasheri M. Evaluating the quality of materials and methods for writings of final proposal in clinical trial studies in Shahrekord University of Medical Sciences based on Consort checklist. *Journal of Clinical Nursing and Midwifery*. 2014; 2(4):1-7.
28. Ahrary Z, Khosravan S, Alami A, Najafi Nesheli M. The effects of a supportive-educational intervention on women with type 2 diabetes and diabetic peripheral neuropathy: a randomized controlled trial. *Clinical Rehabilitation*. 2020; 34(6):794-802. <https://doi.org/10.1177/0269215520914067>
29. Hemmati Maslakkpak M, Shahbaz A, Parizad N, Ghafourifard M. Preventing and managing diabetic foot ulcers: application of Orem's self-care model. *International Journal of Diabetes in Developing Countries*. 2018; 38(2):165-72. <https://doi.org/10.1007/s13410-017-0570-5>
30. Mansouri A, Shahdadi H. The effect of self-care education program based on "Oram Self-care Model" on self-efficacy of Women with gestational diabetes mellitus. *Journal of Diabetes Nursing*. 2017; 5(2):157-66.
31. Baraz S, Zarea K, Shahbazian HB. Impact of the self-care education program on quality of life in patients with type II diabetes. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2017; 11:S1065-S8. <https://doi.org/10.1016/j.dsx.2017.07.043>
32. Borji M, Otaghi M, Kazembeigi S. The impact of Orem's self-care model on the quality of life in patients with type II diabetes. *Biomedical and Pharmacology Journal*. 2017; 10(1):213-20.
33. Ganjloo J, Talebi Z, Asaroudi A, Rakhshani M. Comparative assessment of effect of education in the orem's self care model way with current method on the quality of life of diabetic type 2 patients. *Journal of sabzevar university of medical sciences*. 2015; 2(5):748-57.
34. shahbaz a, hemmati maslakkpak m, nejadrahim r, khalkhali h. The effect of implementing Orem's self-care program on self-care behaviors in patients with diabetic foot ulcer *Nursing and Midwifery Journal*. 2016; 14(2):108-17.
35. Ghafourifard M, Ebrahimi H. The effect of Orem's self-care model-based training on self-care agency in diabetic patients. *Avicenna Journal of Nursing and Midwifery Care*. 2015; 23(1):5-13.
36. Khosravan S, Njafi M, Alami A. Self-care ability of women with diabetes who suffered from peripheral neuropathy and its related needs based on Orem's self-care model. *Medical - Surgical Nursing Journal*. 2015; 4(3):44-51
37. Amirian A, Jandaghian Bidgoli M, Khorvash F, Abdi F, Shaterian N, Jafarimehr M. Investigating the Association of Religious Beliefs with Reproductive Behaviors in Reproductive Aged Women: A Systematic Review. *Journal of Isfahan Medical School*. 2023; 41(737):853-86
38. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *Journal of Pharmacology and pharmacotherapeutics*. 2010; 1(2):100-7. <https://doi.org/10.4103/0976-500X.72352>
39. Karkazloo NV, Daryasari GA, Farahani B, Mohammadnezhad E, Sajjadi A. The study of self-care agency in patients

- with diabetes (Ardabil). *Modern Care Journal*. 2011; 8(4).
40. Kashfi S, Khani JA, Bahadori KR, HAtami M. Evaluation of the effects of educating about nutrition and jogging on the blood sugar of type II diabetic patients of a clinic in Shiraz, Iran. 2009.
 41. Norouzi S, Ghalibaf AK, Sistani S, Banazadeh V, Keykhaei F, Zareishargh P, et al. A mobile application for managing diabetic patients' nutrition: A food recommender system. *Archives of Iranian Medicine*. 2018; 21(10):466.
 42. Lee CS, Wang MH, Acampora G, Hsu CY, Hagraas H. Diet assessment based on type-2 fuzzy ontology and fuzzy markup language. *International Journal of Intelligent Systems*. 2010; 25(12):1187-216. <https://doi.org/10.1002/int.20449>
 43. Kohinor MJ, Stronks K, Nicolaou M, Haafkens JA. Considerations affecting dietary behaviour of immigrants with type 2 diabetes: a qualitative study among Surinamese in the Netherlands. *Ethnicity & Health*. 2011;16(3):245-58. <https://doi.org/10.1080/13557858.2011.563557>
 44. Sanei sistani S, Zadimir M. Effect of Using Self-Care Behaviors in the Diabetic Foot Prevention among Patients with Diabetic Foot Ulcer: A Systematic Review. *Journal of Diabetes Nursing*. 2021; 9(1):1356-66.
 45. Singh S, Jajoo S, Shukla S, Acharya S. Educating patients of diabetes mellitus for diabetic foot care. *Journal of Family Medicine and Primary Care*. 2020; 9(1):367. 10.4103/jfmipc.jfmipc_861_19
 46. Li J, Gu L, Guo Y. An educational intervention on foot self-care behaviour among diabetic retinopathy patients with visual disability and their primary caregivers. *Journal of Clinical Nursing*. 2019; 28(13-14):2506-16. <https://doi.org/10.1111/jocn.14810>
 47. Bolourchifard F, Neishabory M, AbedSaeedi Z. The effects of group and individual education on the level of the knowledge, attitude and practice of self care in patients with diabetic foot ulcer. *Iran Journal of Nursing*. 2009; 22(59):33-41.
 48. Azmi N, Aris M, Nasreen H, Che-Ahmad A. Diabetic foot care practice and its associated factors among type 2 diabetes mellitus patients attending Primary Health Clinics in Kuantan, Malaysia: a cross sectional study. *IJUM Medical Journal Malaysia*. 2020; 19(2). <https://doi.org/10.31436/imjm.v19i2.1560>
 49. Brown DR, Gary LE. Social support network differentials among married and nonmarried Black females. *Psychology of Women Quarterly*. 1985; 9(2):229-41. <https://doi.org/10.1111/j.1471-6402.1985.tb00874.x>
 50. Kooshyar H, Shoorvazi M, Dalir Z, Hoseini SM. Surveying Health-Related Quality of Life and Related Factors in the Diabetic Elderly in Mashhad in 2013. *Journal of Rafsanjan University of Medical Sciences*. 2015; 14(3):175-88.
 51. Salehi L, Salaki S, Alizadeh L. Health-related Quality of Life among Elderly Member of Elderly Centers in Tehran. *Iranian Journal of Epidemiology*. 2012; 8(1):14-20.
 52. Hagedoorn M, Van Yperen NW, Coyne JC, van Jaarsveld CH, Ranchor AV, van Sonderen E, et al. Does marriage protect older people from distress? The role of equity and recency of bereavement. *Psychology and Aging*. 2006; 21(3):611. <https://doi.org/10.1037/0882-7974.21.3.611>
 53. Friedman EM, Love GD, Rosenkranz MA, Urry HL, Davidson RJ, Singer BH, et al. Socioeconomic status predicts objective and subjective sleep quality in aging women. *Psychosomatic medicine*. 2007; 69(7):682-91. 10.1097/PSY.0b013e31814ceada
 54. Taghdisi Heydarian SM, Amiri M, Touzandeh Jani H. The effectiveness of self-control and communication skills on emotional regulation, perceived pain severity and self-care behaviors in diabetic neuropathy. *Journal of Fundamentals of Mental Health*. 2018; 20(6):377-87.
 55. Jacobs M, Snow J, Geraci M, Vythilingam M, Blair R, Charney DS, et al. Association between level of emotional intelligence and severity of anxiety in generalized social phobia. *Journal of Anxiety Disorders*. 2008; 22(8):1487-95. <https://doi.org/10.1016/j.janxdis.2008.03.003>
 56. Cezaretto A, Ferreira SRG, Sharma S, Sadeghirad B, Kolahdooz F. Impact of lifestyle interventions on depressive symptoms in individuals at-risk of, or with, type 2 diabetes mellitus: a systematic review and meta-analysis of randomized controlled trials. *Nutrition, Metabolism and Cardiovascular Diseases*. 2016; 26(8):649-62. <https://doi.org/10.1016/j.numecd.2016.04.009>
 57. Gabbay RA, Añel-Tiangco RM, Dellasega C, Mauger DT, Adelman A, Van Horn DH. Diabetes nurse case management and motivational interviewing for change (DYNAMIC): results of a 2-year randomized controlled pragmatic trial. *Journal of Diabetes*. 2013; 5(3):349-57. <https://doi.org/10.1111/1753-0407.12030>

58. Mirmahdi A, Ansari Jaber A, Shafiepour M, Sayadi A. The Effect of Psychological Training with Progressive Muscle Relaxation on the Stress, Anxiety, and Depression in Diabetes Type 2 Patients. *Community Health Journal*. 2019; 13(4):32-43. 10.22123/CHJ.2020.202309.1363
59. Seligman ME. *Flourish: A visionary new understanding of happiness and well-being*: Simon and Schuster; 2012.
60. Rajabi S, Alimoradi K, Moradi N. The relationship between structural of spiritual well-being and marital satisfaction with the mediating role of emotional intelligence and self-differentiation. *Psychological Methods and Models*. 2017; 8(28):19-48. 20.1001.1.22285516.1396.8.28.2.9
61. adare J, Olamoyegun M, Gbadegehin B. Medication adherence and direct treatment cost among diabetes patients attending a tertiary healthcare facility in Ogbomosho, Nigeria. *Malawi Medical Journal*. 2015;27(2):65-70.
62. Bazargan M, Smith J, Yazdanshenas H, Movassaghi M, Martins D, Orum G. Non-adherence to medication regimens among older African-American adults. *BMC Geriatrics*. 2017; 17(1):163. 10.1186/s12877-017-0558-5
63. Dickson VV, Buck H, Riegel B. Multiple comorbid conditions challenge heart failure self-care by decreasing self-efficacy. *Nursing Research*. 2013; 62(1):2-9. 10.1097/NNR.0b013e31827337b3
64. Tuha A, Getie Faris A, Andualem A, Ahmed Mohammed S. Knowledge and practice on diabetic foot self-care and associated factors among diabetic patients at Dessie referral hospital, northeast Ethiopia: mixed method. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 2021: 1203-14. <https://doi.org/10.2147/DMSO.S300275>
65. Lipscombe C, Smith KJ, Gariépy G, Schmitz N. Gender differences in the relationship between anxiety symptoms and physical inactivity in a community-based sample of adults with type 2 diabetes. *Canadian Journal of Diabetes*. 2014; 38(6):444-50. <https://doi.org/10.1016/j.cjcd.2013.12.002>
66. Khadivzadeh T, Azhari S, Esmaily H, Akhlaghi F, Sardar MA. Effects of self-care education on perceived stress in women with gestational diabetes under insulin treatment. *Evidence Based Care*. 2015; 5(3):7-18.
67. Moradi F, Ghadiri-Anari A, Enjezab B. COVID-19 and self-care strategies for women with gestational diabetes mellitus. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020; 14(5):1535-9. <https://doi.org/10.1016/j.dsx.2020.08.004>
68. Salsali M, Salehi H, Noktehdan H, Hoseini M, KhoshNiyat Nikoo M, Ebrahimi S. Evaluation of the effect of using Symptom Focused Management Model (SFMM) on the amount of HbA1c, knowledge, and self care behavior of type 2 diabetes patients. *Evidence Based Care*. 2013; 3(2):27-36. 10.22038/ebcj.2013.1064
69. Shojaei SF, Masoumi R. The importance of mental health training for psychologists in COVID-19 outbreak. *Middle East Journal of Rehabilitation and Health Studies*. 2020; 7(2). <https://doi.org/10.5812/mejrh.102846>
70. Carolan M, Steele C, Margetts H. Attitudes towards gestational diabetes among a multiethnic cohort in Australia. *Journal of Clinical Nursing*. 2010; 19(17-18):2446-53. <https://doi.org/10.1111/j.1365-2702.2010.03305.x>
71. Dolatian M, Alavi Majd H. Comparison of pregnancy self-care, perceived social support and perceived stress of women with gestational diabetes and healthy pregnant women. *Iranian Journal of Endocrinology and Metabolism*. 2014; 16(3):156-64.
72. Emagholi Khooshehchin T, Keshavarz Z, Afrakhteh M, Shakibazadeh E, Faghihzadeh S. Beliefs, attitudes and coping behaviors of mothers with gestational diabetes: a qualitative study. *Nursing and Midwifery Journal*. 2016; 14(9):791-801.